LIST OF FIGURES

Figure 3.1 S. obtusifolia growing in wild field 32
Figure 3.2 S. obtusifolia growing in
Figure 3.3 S. obtusifolia growing in
Figure 3.4 S. obtusifolia seedlings showing germination in the experimental field 33
Figure 3.5 S. obtusifolia from different provenances in the experimental field 33
Figure 3.6 S. obtusifolia young plants in the experimental field
Figure 3.7 S. obtusifolia in buds, flowers and pods grown in the experimental field 35
Figure 3.8 Mature <i>S. obtusifolia</i> plants in the experimental field
Figure 3.9 Growth patterns of <i>S. obtusifolia</i> of 20 accessions
Figure 3.10 Flowering curve of S. obtusifolia of 20 accessions
Figure 3.11 Fruiting performance of 20 accessions of <i>S. obtusifolia</i>
Figure 3.12 Clustering of <i>S. obtusifolia</i> accessions based on phenological characteristics
Figure 4.1 Matured S. obtusifolia
Figure 4.2 Root of S. obtusifolia 55
Figure 4.3 Leaf of <i>S. obtusifolia</i>
Figure 4.4 Flower of <i>S. obtusifolia</i>
Figure 4.5 Pods of <i>S. obtusifolia</i>
Figure 4.6 Plant heights of <i>S. obtusifolia</i> of 20 accessions
Figure 4.7 Root lengths of <i>S. obtusifolia</i> of 20 accessions

Figure 4.8 Number of branches of <i>S. obtusifolia</i> of 20 accessions	63
Figure 4.9 Internode lengths of <i>S. obtusifolia</i> of 20 accessions	63
Figure 4.10 Girth of <i>S. obtusifolia</i> of 20 accessions	64
Figure 4.11 Leaflet (top) lengths of <i>S. obtusifolia</i> of 20 accessions	64
Figure 4.12 Leaflet (top) widths of <i>S. obtusifolia</i> of 20 accessions	65
Figure 4.13 Rachis with petiole lengths of <i>S. obtusifolia</i> of 20 accessions	65
Figure 4.14 Stipule lengths of <i>S. obtusifolia</i> of 20 accessions	66
Figure 4.15 Stipel lengths of <i>S. obtusifolia</i> of 20 accessions	66
Figure 4.16 Bract lengths of <i>S. obtusifolia</i> of 20 accessions	67
Figure 4.17 Pedicel lengths of <i>S. obtusifolia</i> of 20 accessions	67
Figure 4.18 Sepal lengths of <i>S. obtusifolia</i> of 20 accessions	68
Figure 4.19 Petal lengths of <i>S. obtusifolia</i> of 20 accessions	68
Figure 4.20 Anther lengths of <i>S. obtusifolia</i> of 20 accessions	69
Figure 4.21 Filament lengths of <i>S. obtusifolia</i> of 20 accessions	69
Figure 4.22 Pistil lengths of <i>S. obtusifolia</i> of 20 accessions	70
Figure 4.23 Number of Flowers per Plant of <i>S. obtusifolia</i> of 20 accessions	70
Figure 4.24 Number of Pods per Plant of <i>S. obtusifolia</i> of 20 accessions	71
Figure 4.25 Pod lengths of <i>S. obtusifolia</i> of 20 accessions	71
Figure 4.26 Number of seeds per pod of <i>S. obtusifolia</i> of 20 accessions	72
Figure 4.27 Seed weights of <i>S. obtusifolia</i> of 20 accessions	72
Figure 4.28 Correlation plot of morphological characteristics of S. obtusifolia	77

Figure 4.29 Principal components and their cumulative proportion of variance 81
Figure 4.30 Classification of <i>S. obtusifolia</i> based on morphological characteristics. 83
Figure 5.1 Metaphase karyotype plate and ideogram from the germinated root tip of the accession ASN (Asansol)
Figure 5.2 Metaphase karyotype plate and ideogram from the germinated root tip of the accession BDG (Bandhavgarh)
Figure 5.3 Metaphase karyotype plate and ideogram from the germinated root tip of the accession BNP (Bishnupur)
Figure 5.4 Metaphase karyotype plate and ideogram from the germinated root tip of the accession BPR (Bolpur)
Figure 5.5 Metaphase karyotype plate and ideogram from the germinated root tip of the accession DHD (Dahod)
Figure 5.6 Metaphase karyotype plate and ideogram from the germinated root tip of the accession DHN (Dehradun) 100
Figure 5.7 Metaphase karyotype plate and ideogram from the germinated root tip of the accession DNG (Devendranagar)
Figure 5.8 Metaphase karyotype plate and ideogram from the germinated root tip of the accession GBP (Gobindapur)
Figure 5.9 Metaphase karyotype plate and ideogram from the germinated root tip of the accession HDR (Haridwar)
Figure 5.10 Metaphase karyotype plate and ideogram from the germinated root tip of the accession JBP (Jabalpur)
Figure 5.11 Metaphase karyotype plate and ideogram from the germinated root tip of the accession JPR (Jaipur)
Figure 5.12 Metaphase karyotype plate and ideogram from the germinated root tip of the accession JRM (Jhargram)

Figure 5.13 Metaphase karyotype plate and ideogram from the germinated root tip of
the accession KLN (Kalyani)107
Figure 5.14 Metanbase karvotype plate and ideogram from the germinated root tip of
the accession KDD (Kharagnur)
the accession KFK (Kharagpur)108
Figure 5.15 Metaphase karyotype plate and ideogram from the germinated root tip of
the accession LKP (Lakshmikantapur)109
Figure 5.16 Metaphase karyotype plate and ideogram from the germinated root tip of
the accession MGL (Mangalore)110
Figure 5.17 Metaphase karvotype plate and ideogram from the germinated root tip of
the accession NGP (Nagpur)
Figure 5.18 Metaphase karyotype plate and ideogram from the germinated root tip of
the accession PRL (Purulia)112
Figure 5.19 Metaphase karyotype plate and ideogram from the germinated root tip of
the accession RPR (Raipur)
Figure 5.20 Metaphase karyotype plate and ideogram from the germinated root tip of
the accession TNG (Tatanagar)
Figure 5.21 Polyploidy of S. obtusifolia 117
Figure 5.22 Variations asymptotic of S. altraifalia of twanty appagaions 110
Figure 5.22 Karyotype asymmetry of 5. obtustiona of twenty accessions
Figure 6.1 Polyketide pathway for the biosynthesis of anthraquinones 124
Figure 6.2 Pathway of rhein 126
Figure 6.3 Molecular structure of Rhein 127
1 gure die Monecular Structure of Ferenin
Figure 6.4 Chromatogram of standard rhein solution of concentration of 5 $\mu\text{g/ml}$. 136
Figure 6.5 Chromatogram of standard rhein solution of concentration of 10 μ g/ml 136
Figure 6.6 Chromatogram of standard rhein solution of concentration of 20 μ g/ml 137

Figure 6.7 Chromatogram of standard rhein solution of concentration of 40 μ g/ml 137
Figure 6.8 Calibration curve generated from chromatogram of standard rhein solution
Figure 6.9 Chromatogram of extracted rhein solution from the leaves of the plants of the accession ASN (Asansol)
Figure 6.10 Chromatogram of extracted rhein solution from the leaves of the plants of the accession BDG (Bandhavgarh)
Figure 6.11 Chromatogram of extracted rhein solution from the leaves of the plants of the accession BNP (Bishnupur)
Figure 6.12 Chromatogram of extracted rhein solution from the leaves of the plants of the accession BPR (Bolpur)
Figure 6.13 Chromatogram of extracted rhein solution from the leaves of the plants of the accession DHD (Dahod)
Figure 6.14 Chromatogram of extracted rhein solution from the leaves of the plants of the accession DHN (Dehradun)
Figure 6.15 Chromatogram of extracted rhein solution from the leaves of the plants of the accession DNG (Devendranagar)
Figure 6.16 Chromatogram of extracted rhein solution from the leaves of the plants of the accession GBP (Gobindapur)
Figure 6.17 Chromatogram of extracted rhein solution from the leaves of the plants of the accession HDR (Haridwar)
Figure 6.18 Chromatogram of extracted rhein solution from the leaves of the plants of the accession JBP (Jabalpur)
Figure 6.19 Chromatogram of extracted rhein solution from the leaves of the plants of the accession JPR (Jaipur)

Figure 6.20 Chromatogram of extracted rhein solution from the leaves of the plants of
the accession JRM (Jhargram) 145
Figure 6.21 Chromatogram of extracted rhein solution from the leaves of the plants of
the accession KLN (Kalyani)146
Figure 6.22 Chromatogram of extracted rhein solution from the leaves of the plants of
the accession KPR (Kharagpur)146
Figure 6.23 Chromatogram of extracted rhein solution from the leaves of the plants of
the accession (Lakshmikantapur)147
Figure 6.24 Chromatogram of extracted rhein solution from the leaves of the plants of
the accession MGL (Mangalore)147
Figure 6.25 Chromatogram of extracted rhein solution from the leaves of the plants of
the accession NGP (Nagpur)148
Figure 6.26 Chromatogram of extracted rhein solution from the leaves of the plants of
the accession PRL(Purulia)148
Figure 6.27 Chromatogram of extracted rhein solution from the leaves of the plants of
the accession RPR (Raipur)149
Figure 6.28 Chromatogram of extracted rhein solution from the leaves of the plants of
the accession TNG (Tatanagar)149
Figure 6.29 Chart for extracted rhein quantity in microgram per gram of dried leaf
against the accessions
Figure 6.30 Clustering of S. obtusifolia of 20 accessions based on the quantity of rhein
extracted from dried leaf154
Figure 6.31 ESI MS spectrum of <i>S. obtusifolia</i> sample solution 157
Figure 7.1 S. obtusifolia Genomic DNA quantification on agarose gel 163
Figure 7.2 Pre-amplification of adaptor ligated libraries

Figure 7.3 DNA Fingerprint (AFLP Primer Pair E-AAC/M-CAG) 169
Figure 7.4 DNA Fingerprint (AFLP Primer Pair E-AAC/M-CTA) 170
Figure 7.5 DNA Fingerprint (AFLP Primer Pair E-ACA/M-CAG) 171
Figure 7.6 DNA Fingerprint (AFLP Primer Pair E-ACA/M-CTA) 172
Figure 7.7 DNA Fingerprint (AFLP Primer Pair E-ACT/M-CAG) 173
Figure 7.8 DNA Fingerprint (AFLP Primer Pair E-ACT/M-CTA) 174
Figure 7.9 DNA Fingerprint (AFLP Primer Pair E-AG/M-CAG) 175
Figure 7.10 DNA Fingerprint (AFLP Primer Pair E-AG/M-CTA) 176
Figure 7.11 DNA Fingerprint (AFLP Primer Pair E-AT/M-CAG) 177
Figure 7.12 DNA Fingerprint (AFLP Primer Pair E-AT/M-CTA) 178
Figure 7.13 UPGMA dendrogram showing the genetic diversities of AFLP fingerprints of 20 accessions
Figure 8.1 Total haploid chromosome length of <i>S. obtusifolia</i> of twenty accessions
Figure 8.2 Overall dendrogram of phenetic analysis of 20 S. obtusifolia accessions