

LIST OF PLATES

| Plate No. | Plate and Diagram captions | Page Nos. |
|-----------|--|-----------|
| 4.1.1 | Histology of testis 400X (Hematoxyline-Eosin Stain). Representative microphotographs of A) Vehicle treated control and B) CPA treated rat where the size of seminiferous tubule in CPA treated rat is remarkably smaller than the vehicle treated control. | |
| 4.2.1 | Histology of testis 400X (Hematoxyline-Eosin Stain). Representative microphotographs of A) vehicle treated control, B) CPA treated rat, C) 0.75 mg lycopene treated, D) 1.5 mg lycopene treated, E) 3.0 mg lycopene treated, F) 4.5 mg lycopene treated rat. Recovery in the seminiferous tubular diameter after treatment with different doses of lycopene to the CPA treated rat which focus the rectification in the sperm density after lycopene treatment | |

LIST4.4OF TABLES

| Table No. | Table captions | Page Nos. |
|-----------|---|-----------|
| 4.1.1 | Effect of cyproterone acetate on body weight and reproductive organo-somatic indices | |
| 4.1.2 | Spermiological sensors of the rat after treatment with CPA | |
| 4.2.1 | Dose dependent corrective efficacy of lycopene on spermatogenic profile in CPA induced infertile rats | |
| 4.3.1 | Effect of lycopene in 15 days, 30 days and 45 days duration dependent group on body weight and reproductive organosomatic indices in CPA induced male rat | |
| 4.3.2 | Toxicity profile after the administration of lycopene for 15 days, 30 days and 45 days to the CPA treated infertile male rats | |
| 4.3.3 | Spematogenic profile after the administration of lycopene for 15 days, 30 days and 45 days to the CPA treated infertile male rats | |
| 4.3.4 | Recovery in the spermatogenic cycle after administration of lycopene for 15 days, 30 days and 45 days | |
| 4.4.1 | Spematogenic profile after cessation of lycopene treatment for | |

| | | |
|--------------|---|--|
| | 15 days, 30 days, 45 days to the CPA treatment male albino rat | |
| 4.4.2 | Levels of serum tryglyceride, VLDL, LDL, HDL after cessation of lycopene treatment for 15 days, 30 days, 45 days to the CPA treated male albino rat | |
| 4.7.1 | Effect of lycopene treatment on serum tryglyceride, VLDL, LDL, HDL profile in albino rat | |
| 4.8.1 | Quantification of the implantation site in both of the uterine horn of the female albino rats mated with the vehicle treated control, CPA treated group and CPA+ lycopene treated rat. | |