

List of Figures

Figure No	Legends	Page No
4.1.	Schematic graphic of study design of present research work.	29
4.2.	Schematic diagram of study design of present research work.	30
5.1.	Represents the pattern of UV-visible spectroscopy absorbance of CCPS and As ^{III} + CCPS association. (a) Indicates the UV-visible spectrogram absorbance of CCPS and (b) Indicates As ^{III} + CCPS association.	34
5.2.	Represents the Fourier transforms infrared spectrum (FTIR) of CCPS and As ^{III} + CCPS. (a) Indicates the FTIR spectrum of CCPS and (b) Indicates the FTIR spectrum of As ^{III} + CCPS association.	35
5.3.	Represents the X-ray diffraction spectroscopy patterns of CCPS and As ^{III} + CCPS association. (a) Indicates the XRD pattern of CCPS, and (b) Indicates the XRD pattern of As ^{III} + CCPS.	35
5.4.	Represents the NMR spectrum patterns of CCPS and As ^{III} + CCPS association (a) Indicates the ¹ H NMR pattern of CCPS and (b) Indicates ¹ H NMR pattern of As ^{III} + CCPS association.	35
5.5.	Represent the electron microscopic study of CCPS and As ^{III} + CCPS association. (A) Indicates the scanning electron microscopy (SEM) of CCPS, (B) Indicates SEM of As ^{III} + CCPS association, (C) Indicates the transmission electron microscopy (TEM) of As ^{III} + CCPS association and (D) Indicates the Diffraction image of As ^{III} + CCPS association.	36
6.1.	(A & B) represents the hepatic SOD and catalase expression on a native gel. Lane division: Lane 1 indicates the control group; Lane 2 as 0.2 ppm	44

	of As ^{III} ; Lane 3 as 0.4 ppm of As ^{III} ; Lane 4 as 0.6 ppm of As ^{III} ; Lane 5 as 0.8 ppm of As ^{III} ; Lane 6 as CCPS; Lane 7 as Cur; Lane 8 as As ^{III} + CCPS and Lane 9 as As ^{III} + Cur.	
6.2.	Protective effects of curcumin and CCPS on SOD and catalase level in liver tissue slices against arsenic exposed rats in duration dependent fashion. Data represent mean \pm SE, N = 6. Presented values here are expressed by one-way ANOVA with Dunnett's Post Hoc t-test *, **, *** indicates p<0.05, p<0.01, p<0.001 versus control group with vehicle, whereas #, ##, ### indicate p<0.05, p<0.01 and p<0.001 versus As ^{III} treatment.	46
6.3.	(A-H) indicates the SOD, catalase and Px expression in liver tissue on polyacrylamide gel. Lane division: Lane 1 indicates control group with vehicle; Lane 2 as As ^{III} ; Lane 3 as H ₂ O ₂ ; Lane 4 as As ^{III} + H ₂ O ₂ ; Lane 5 as As ^{III} + H ₂ O ₂ + Cur; Lane 6 as As ^{III} + H ₂ O ₂ + CCPS; Lane 7, 8 as As ^{III} + H ₂ O ₂ + Cur + CCPS. Fig 2 denotes (D-H) the liver catalase and peroxidase expression on polyacrylamide gel. Lane division is following manner: Lane 1 is control group; Lane 2 as As ^{III} ; Lane 3 as H ₂ O ₂ ; Lane 4 as As ^{III} + H ₂ O ₂ ; Lane 5 as As ^{III} + H ₂ O ₂ + Cur; Lane 6 as As ^{III} + H ₂ O ₂ + CCPS; Lane 7-9 as As ^{III} + H ₂ O ₂ + Cur + CCPS.	47
6.4	(A & B) denotes the activity of liver LDH. Lane division shown in following manner: Lane 1 is control group with vehicle; Lane 2 as As ^{III} ; Lane 3 as As ^{III} + H ₂ O ₂ ; Lane 4 as As ^{III} + CCPS; Lane 5 as As ^{III} + Cur; Lane 6 as As ^{III} + H ₂ O ₂ + CCPS; Lane 7 as As ^{III} + H ₂ O ₂ + Cur; Lane 8 as As ^{III} + H ₂ O ₂ + Cur + CCPS.	48

6.5.	The effects of curcumin and CCPS or combination on DNA fragmentation against arsenic and H ₂ O ₂ induced changes in hepatic slices. Lane division; Lane 1 indicates the control group with vehicle; Lane 2 as As ^{III} ; Lane 3 as H ₂ O ₂ ; Lane 4 as As ^{III} + H ₂ O ₂ ; Lane 5 as As ^{III} + Cur; Lane 6 as As ^{III} + CCPS; Lane 7 as As ^{III} + H ₂ O ₂ + Cur; Lane 8 as As ^{III} + H ₂ O ₂ + CCPS.	48
6.6.	(A & B) The effects of CCPS and curcumin in liver cells on the single cell assay against arsenic and H ₂ O ₂ induced changes in duration dependent fashion. Lane allotment; Lane A is control group with vehicle; Lane B as As ^{III} ; Lane C as H ₂ O ₂ ; Lane D as As ^{III} + H ₂ O ₂ ; Lane E as As ^{III} + Cur; Lane F as As ^{III} + CCPS; Lane G as As ^{III} + H ₂ O ₂ + Cur; and Lane H as As ^{III} + Cur + CCPS + H ₂ O ₂ .	49
6.7.	Schematic diagram denotes the hypothetical mechanism action of curcumin and CCPS against arsenic/H ₂ O ₂ induced liver toxicity. Black colour (+), (-) sign and black line represents the stimulatory and as well as inhibitory effect of arsenic/H ₂ O ₂ respectively. Red colour (+), (-) sign and red line indicates the stimulatory and inhibitory effect of curcumin and CCPS respectively.	55
7.1.	Effect of three doses of Cur on the pattern of estrous cycle against arsenic mediated different groups. Data represent mean ± SE, N = 6, Presented values here are expressed using one-way ANOVA with Dunnett's post Hoc t-test * indicate p<0.05 versus the control with vehicle group, whereas # and ## indicate p<0.05 and p<0.01 versus As ^{III} treatment.	61
7.2.	The spectrophotometry (A & B) data showing the effects of three doses of curcumin on uterine endogenous antioxidants (SOD & catalase) levels. Electrozymogram (C & D) images also showing the uterine endogenous	63

	enzymatic antioxidant expression. E & F data shows the band density (%) of SOD and catalase expression respectively. Data represent mean \pm SE, N = 6, Presented values here are expressed using one-way ANOVA with Dunnett's post Hoc t-test *, *** indicate $p < 0.05$, $p < 0.001$ versus vehicle with control, whereas #, ### indicate $p < 0.05$, $p < 0.001$ versus As ^{III} treatment.	
7.3.	A & B denotes the effects of different doses of curcumin on ovarian key regulatory steroidogenic enzyme activities in arsenic treated animals. C, D & E data shows that the effects of different doses of curcumin on gonadotropin and estradiol hormones in arsenicated rats. All data represent mean \pm SE, N = 6, Presented values here are expressed using one-way ANOVA with Dunnett's post Hoc t-test *, *** indicate $p < 0.05$, $p < 0.001$ versus vehicle with control, whereas # indicate $p < 0.05$ versus As ^{III} treatment.	64
7.4.	Schematic diagram showing the hypothetical mechanism action of curcumin against arsenic-induced uterine toxicity. Red colour (+), (-) sign and black line expresses the stimulatory and as well as the inhibitory effect of arsenic respectively. Black colour (+), (-) sign and green line also expresses the stimulatory and inhibitory effects of curcumin respectively.	69
8.1.	Effects of three doses of CCPS on the pattern of estrous cycle against arsenic ingested groups. The all data here represent mean \pm SE, N = 6. Presented values here are expressed using one-way ANOVA with Dunnett's post Hoc t-test * indicate $p < 0.05$ versus control with vehicle, whereas #, ## and ### indicate $p < 0.05$, $p < 0.01$ and $p < 0.001$ versus As ^{III} treatment.	74

8.2.	The spectrophotometry (A & B) data showing the effects of three doses of CCPS on uterine endogenous enzymes antioxidant (SOD & catalase) activities. Electrozymogram (C & D) images showing the effects of different doses of CCPS on uterine endogenous enzymes antioxidant expression. E & F data shows the band density (%) of SOD and catalase respectively. Here, data represent mean \pm SE, N = 6, Presented values here are expressed using one-way ANOVA with Dunnett's post Hoc t-test *, ***, indicates $p < 0.05$, $p < 0.001$ versus the control with vehicle, whereas #, ##, ### indicates, $p < 0.05$, $p < 0.01$, $p < 0.001$ versus As^{III} treatment.	76
8.3.	A & B denotes the effects of three doses of CCPS on ovarian key regulatory steroidogenic enzyme activity. C, D & E data shows that effects on gonadotrophin and estradiol hormones. Data are represent mean \pm SE, N = 6, Presented values here are expressed using one-way ANOVA with Dunnett's post Hoc t-test *, **, *** indicate $p < 0.05$, $p < 0.01$, $p < 0.001$ versus the vehicle with control, whereas ##, ### indicate $p < 0.01$, $p < 0.001$ versus As^{III} .	77
8.4.	Schematic figure showing the hypothetical mechanism of action of CCPS against arsenic induced uterine toxicity. Red colour (+) and (-) sign expresses stimulatory and inhibitory effect of arsenic respectively. Black colour (+) and (-) sign expresses the stimulatory and also the inhibitory effect of CCPS respectively.	81
9.1.	Protective effects of curcumin and CCPS on the pattern of estrous cycle against arsenic ingested different group of rats. Bar diagram represent mean \pm SE, N = 6. Presented values here are expressed using one-way ANOVA with Dunnett's post Hoc t-test * indicate $p < 0.05$ versus the	88

	control with vehicle, whereas # and ## indicate $p < 0.05$, $p < 0.01$ versus As^{III} treatment.	
9.2.	Protective effects of curcumin and CCPS on MDA and CD in uterine tissue against the arsenic challenged rats. Bar diagram represent mean \pm SE, N = 6. Presented values here are expressed using one-way ANOVA with Dunnett's post Hoc t-test **,*** indicate $p < 0.01$ and $p < 0.001$ versus the control with vehicle, whereas # and ### indicate $p < 0.05$, and $p < 0.001$ versus As^{III} treatment.	89
9.3.	(A, B & C) Spectrophotometric analysis showed the protective effects of curcumin and CCPS on uterine endogenous antioxidant enzymes activities against arsenic ingested different group of rats. Electrozymogram showed the protective effects of curcumin and CCPS on uterine expression of antioxidant enzymes. Table showing: (D) Band density (%) of SOD, catalase and PX respectively. Bar diagram represent mean \pm SE, N = 6. Presented values here are expressed using one-way ANOVA with Dunnett's post Hoc t-test *, *** indicate $p < 0.05$ and $p < 0.001$ versus the control with vehicle, whereas # and ### indicate $p < 0.05$ and $p < 0.001$ versus As^{III} treatment.	90
9.4.	(A) Spectrophotometric measurement showed the protective effects of curcumin and CCPS on uterine tissue necrosis level (LDH) against arsenic ingested different group of rats. (B) Electrozymogram showed that the protective effects of curcumin and CCPS on uterine tissue necrosis activity (LDH). Figure showing: (C) Band density (%) of serum LDH status. Data represent mean \pm SE, N = 6. Presented values here are expressed using one-way ANOVA with Dunnett's post Hoc t-test *,**,	91

	*** indicate $p < 0.05$, $p < 0.01$ and $p < 0.001$ versus the control with vehicle, whereas ##, ### indicate $p < 0.01$ and $p < 0.001$ versus As^{III} treatment.	
9.5.	(A & B) Protective effects of curcumin and CCPS on uterine DNA damages against arsenic ingested different group of rats. Uterine DNA was evaluated by image J software. Figure B is showing the effect of curcumin and CCPS on DNA comet in uterine cells. Lane distribution; Lane B ₁ indicate control with vehicle; Lane B ₂ as As^{III} ; Lane B ₃ as Cur; Lane B ₄ as As^{III} + Cur; Lane B ₅ as CCPS; Lane B ₆ as As^{III} + CCPS; Lane B ₇ as As^{III} + Cur + CCPS.	92
9.6.	(A, B and C) Protective effects of curcumin and CCPS on serum vitamins and Hcy levels against arsenic ingested rats. Data represent mean \pm SE, N = 6. Presented values here are expressed using one-way ANOVA with Dunnett's post Hoc t-test *, **, *** indicates $p < 0.05$, $p < 0.01$ and $p < 0.001$ versus the control with vehicle, whereas #, ## and ### indicate $p < 0.05$, $p < 0.01$, $p < 0.001$ versus As^{III} treatment.	93
9.7.	(A & B) Protective effects of curcumin and CCPS on ovarian steroidogenesis and gonadotrophins against arsenic ingested different group of rats. Data represent mean \pm SE, N = 6. Presented values here are expressed using one-way ANOVA with Dunnett's post Hoc t-test *, **, *** indicate $p < 0.05$, $p < 0.01$ and $p < 0.001$ versus the control with vehicle, whereas #, ##, ### indicate $p < 0.05$, $p < 0.01$ and $p < 0.001$ versus As^{III} treatment.	94
9.8.	(A, B, C & D) Protective effects of curcumin and CCPS on pro-inflammatory markers against arsenic challenged group. All data represent mean \pm SE, N = 6. Presented values here are expressed using	95

	<p>one-way ANOVA with Dunnett's post Hoc t-test *, **, *** indicate p<0.05, p<0.01 and p<0.001 versus the control with vehicle, whereas #, ## ### indicate p<0.05, p<0.01 and p<0.001 versus As^{III} treatment.</p>	
9.9A and 9B.	<p>Represent the protective effects of curcumin and CCPS in uterine and ovarian cell morphology against arsenic ingested different group of rats. Fig. 9A shows the uterine cell morphology. Here PM, MM EM denotes Perimetrium, Myometrium and Endometrium of the uterine layers and SG denotes the secretory glands. Arsenic treatment rats degenerates these layers but CCPS treatment rat recovered the three uterine layers. Fig. 9B shows the ovarian cell morphology. Here, AF denotes the atretic follicle. Numbers of atretic follicles here increased in arsenicated group.</p>	96
9.10.	<p>Schematic scheme represent the hypothetical mechanism of curcumin and CCPS alone or combination action against arsenic caused uterine toxicity. Black colour (+), (-) sign and green line indicate stimulatory and inhibitory effects of arsenic respectively. Black colour (+), (-) sign and black line indicates stimulatory and inhibitory effects of curcumin and CCPS respectively.</p>	103
10.1.	<p>Represents the curative effects of CCPS on estrous cycle patterns. Scale bar in the cytological assessment is represented at 50 mm.</p>	110
10.2A.	<p>Represents the curative effect of CCPS in arsenicated rats on liver markers. These data were expressed as means \pm SE, N=6 using one way ANOVA in association with Dunnett's post Hoc t-test. Significant variations of data were expressed at *p<0.05, and ***p<0.001 control group with vehicle and at #p<0.05, and ###p<0.001 with the arsenic challenged group.</p>	111

10.2B.	Represents the curative effect of CCPS in arsenicated rats on the lipid profiles. These data were represented as means \pm SE, N=6 using one way ANOVA in association with Dunnett's post Hoc t-test. Significant variations of data were expressed at *p<0.05, **p<0.01 and ***p<0.001 control group with vehicle and at ##p<0.01, and ###p<0.001 with the arsenic challenged group.	112
10.3.	Represents the curative effects of CCPS on the uterine and ovarian oxidative stress markers against arsenic-treated rats by spectrophotometric evaluation. These results were represented as mean \pm SE, N=6 using one way ANOVA in association with Dunnett's post Hoc t-test. Significant variations of data were expressed at *p<0.05, **p<0.01 and ***p<0.001 when control group with vehicle and at #p<0.05, ##p<0.01 and ###p<0.001 with the arsenic challenged group.	113
10.4A	Represents the curative effects of CCPS on the uterine and ovarian antioxidant enzymes activity against arsenic-treated rats when evaluated spectrophotometrically. These results were represented as means \pm SE, N=6 using one way ANOVA in association with Dunnett's post Hoc t-test. Significant variations of data were expressed at the level of *p<0.05, **p<0.01 and ***p<0.001 when compared control with vehicle group and #p<0.05, and ###p<0.001 when compared with arsenic challenged group.	114
10.4B.	CCPS showed a remedial effect on uterine and ovarian antioxidant enzymes (SOD, catalase and GPx) activities against arsenic treated rats. These results were represented as mean \pm SE, N=6 using one way ANOVA in association with Dunnett's post Hoc t-test. Significant variations of data were expressed at the level of *p<0.05, and ***p<0.001	115

	when compared control group with vehicle whereas at #p<0.05, and ###p<0.001 with arsenic challenged group.	
10.5.	Spectrophotometric analysis denoted that CCPS improved the LDH status (5A) in arsenic treated rats. Zymographic data also focused that CCPS restored the serum LDH towards normalcy (5B). Data here represented as mean ± SE, N=6 using one way ANOVA in association with Dunnett's post Hoc t-test. Significant variations of data were expressed at the level of *p<0.05 and ***p<0.001 compared control group with vehicle and at ###p<0.001 with the arsenic challenged group.	116
10.6.	Represents the remedial effect of CCPS on circulating level of vitamins and Hcy in arsenic-treated rats. These results were represented as mean ± SE, N=6 using one way ANOVA in association with Dunnett's post Hoc t-test. Significant variations of data were expressed at **p<0.01 and ***p<0.001 which compared the control with vehicle and at ###p<0.001 with arsenic challenged group.	117
10.7.	Represents the corrective effects of CCPS on the hormonal level, ovarian steroidogenic key enzymatic activities, and the uterine Esr-1 in arsenicated rats. These results were represented as mean ± SE, N=6 using one way ANOVA in association with Dunnett's post Hoc t-test. Significant variations of data were expressed at *p<0.05, and ***p<0.001 level, compared the control with vehicle and at #p<0.05, ##p<0.01 and ###p<0.001 with arsenic challenged group.	118
10.8.	Shows the therapeutic efficacy curative of CCPS on the status inflammatory markers against arsenication in rats. These results were represented as mean ± SE, N=6 using one way ANOVA in association	119

	<p>with Dunnett's post Hoc t-test. Significant variations of data were expressed at the level of ***$p < 0.001$ compared the control with vehicle and #$p < 0.05$, ##$p < 0.01$ and ###$p < 0.001$ with arsenic challenged group.</p>	
10.9.	<p>The revival mode CCPS action successfully improved the apoptotic protein expression on arsenic- affected rats. These results were represented as mean \pm SE, N=6 using one way ANOVA in association with Dunnett's post Hoc t-test. Significant variations of data were expressed at the level of *$p < 0.05$, **$p < 0.01$ and ***$p < 0.001$ compared the control with vehicle and at #$p < 0.05$, ##$p < 0.01$, and ###$p < 0.001$ with arsenic challenged group.</p>	120
10.10.	<p>The curative mode of CCPS successfully recovered the gene expression in the arsenic- affected rats. These results were represented as mean \pm SE, N=6 using one way ANOVA in association with Dunnett's post Hoc t-test. Significant variations of data were expressed at the level of *$p < 0.05$ and ***$p < 0.001$ compared control group with vehicle and at #$p < 0.05$, ##$p < 0.01$, and ###$p < 0.001$ with the arsenic challenged group.</p>	121
10.11.	<p>Represents the therapeutic potential of CCPS on the sex organs histo morphology in arsenic-ingested rats. Uterine cell morphology revealed the loss of uterine layers (PM- Perimetrium, MM- Myometrium and, EM- Endometrium) along with the loss of secretory glands (SG). Dietary CCPS played a crucial role in rejuvenating the uterine layers along with the secretory glands. In ovarian cell morphology showed that, the numbers of atretic follicles were increased following arsenication. CCPS treatment diminished the numbers of ovarian follicular artesia at a considerable level.</p>	122

10.12.	CCPS successfully alleviates the arsenic affected female infertility status. CCPS also improved pregnancy outcome. These results were represented as mean \pm SE, N=6 using one way ANOVA in association with Dunnett's post Hoc t-test. Significant variations of data were expressed at the level of * $p < 0.05$ and *** $p < 0.001$ compared control group with vehicle and at # $p < 0.05$ and ### $p < 0.001$ with the arsenic challenged group.	123
10.13.	Represents mechanism of actions of CCPS against arsenic mediated female repro-toxicity. Here, (+) sign indicate the stimulatory effect (-) sign also indicate inhibitory effect.	130
11.1.	Represents the solubility of curcumin and encapsulated chitosan nanoparticles (ECNPs).	132
11.2.	Represents the FTIR spectrum of (a) curcumin, (b) chitosan, and (c) ECNPs.	133
11.3.	Represents the size characterization of the ECNPs using TEM studies.	133
11.4.	Analysis of (a) curcumin, (b) chitosan and (c) ECNPs using the help of X-ray diffraction spectroscopic.	134
12.1.	Effect of different doses of ECNPs on the pattern of estrous cycle against arsenicated mediated different group of rats. Data represent mean \pm SE, N = 6. Presented values here are expressed using one way ANOVA in association with Dunnett's post Hoc t-test. * indicates $p < 0.05$ versus the control with vehicle group, whereas # and ## indicate $p < 0.05$ and $p < 0.01$ versus As ^{III} treatment.	142
12.2.	The spectrophotometric (A & B) data showing the effects of different doses of ECNPs on uterine endogenous antioxidant enzymes activities. Electrozymogram (C & D) images showing the effects of different doses	144

	<p>of ECNPs on uterine endogenous antioxidant enzymes expression. E & F shows the band density (%) of SOD and catalase respectively. Data represent mean \pm SE, N = 6. Presented values here are expressed using one way ANOVA in association with Dunnett's post Hoc t-test. *, *** indicate $p < 0.05$, $p < 0.001$ versus the control with vehicle group, whereas #, ##, ### indicate $p < 0.05$, $p < 0.01$ and $p < 0.001$ versus As^{III} treatment.</p>	
12.3.	<p>A & B denote the effects of different doses of ECNPs on ovarian key regulatory steroidogenic enzyme activities. C, D & E show the effects of different doses of ECNPs on gonadotrophin and estradiol hormones. Data represent mean \pm SE, N = 6. Presented values here are expressed using one way ANOVA in association with Dunnett's post Hoc t-test. *, *** indicates $p < 0.05$, $p < 0.001$ versus the control with vehicle group, whereas #, ##, ### indicate $p < 0.05$, $p < 0.01$ and $p < 0.001$ versus As^{III} treatment.</p>	145
12.4.	<p>Schematic diagram showing the hypothetical mechanism action of curcumin against arsenic induced uterine toxicity. Black colour (+), (-) sign showing stimulatory and inhibitory effect of arsenic respectively. Red colour (+), (-) sign showing stimulatory and inhibitory effects of ECNPs respectively.</p>	149