

List of figures

Experiment	Serial no.	Figure No.	Title	Page no.
	1.	Figure 2.1	Arsenic affected area in West Bengal	9
	2.	Figure 2.2	Schematic diagram of the pathway of inorganic arsenic metabolism in liver	14
	3.	Figure 2.3	Phases of Wistar Rat's Estrous Cycle A: Proestrous B: Estrous C: Metestrous D: Diestrous	34
	4.	Figure 2.4	Vitamin B ₁₂	41
	5.	Figure 2.5	Role of Vitamin B ₁₂ and Folic acid in DNA Synthesis	46
	6.	Figure 3	A schematic diagram of the present experimental deigned	64
Experiment-I	7.	Figure 5.1.1	Dose dependent changes of catalase and superoxide dismutase activities in uterine and ovarian tissue after arsenic treatment. Data represents here is mean \pm SE, N=6, p<0.001. ANOVA followed by multiple student's t test. Values of bar diagrams with same superscript did not differ from each other significantly.	70
	8.	Figure 5.1.2	Dose dependent changes of peroxidase activities in uterine tissue after arsenic treatment in 8% native polyacrylamide gel electrophoresis.	71
	9.	Figure 5.1.3	Dose dependent changes of conjugated diene and malondialdehyde levels in uterine and ovarian tissue after arsenic treatment. Data represents here is mean \pm SE, N=6, p<0.001 and ANOVA followed by multiple student's t test. Values of bar diagrams with same superscript did not differ from each other significantly.	71

	10.	Figure 5.1.4	Histological sections show the ovaries of (A) vehical treated control focusing normal folicals. (B) after sodium arsenite treated (0.2 ppm) rat. (C) after sodium arsenite treated(0.4ppm) rat showing follicular atresia	72
	11	Figure 5.1.5	Histological sections show the uterus of (A) vehical treated control focusing normal internal lumem diameter, (B) No change internal lumem dameter after sodium arsenite treated (0.2ppm) rat. (C) after sodium arsenite treated (0.4 ppm) rat showing reduce of internal lumen diameter.	72
Experiment II	12	Figure 5.2.1	Duration dependent changes of catalase and superoxide dismutase activities in uterine and ovarian tissue after arsenic treatment. Data represents here is mean \pm SE, N=6, $p < 0.001$ and ANOVA followed by multiple student's t test. Values of bar diagrams with same superscript did not differ from each other significantly	79
	13	Figure 5.2.2	Duration dependent changes of conjugated diene and malondialdehyde levels in uterine tissue after arsenic treatment. Data represents here is mean \pm SE, N=6, $p < 0.001$ and ANOVA followed by multiple student's t test. Values of bar diagrams with same superscript did not differ from each other significantly.	79
	14	Figure 5.2.3	Histological sections show the ovaries of (A) vehical treated control focusing normal folicals, (B) after sodium arsenite treated(0.4ppm) rat for 16 days focusing normal folicals, (D) vehical treated control focusing normal folicals for 28 days, (E) after sodium arsenite treated (0.4ppm) rat for 28 days showing follicular atresia.	80
Experiment- III	15	Figure 5.3.1	Correction of ovarian $\Delta^5,3\beta$ -HSD and 17β -HSD in arsenic induced toxicity with co-administration of Vit-B ₁₂ (0.07 μ g) and/ or	87

folic acid (4.0 μ g). Each bar represents mean \pm SE, N=6. ANOVA followed by two tailed student's t test were used to find out statistical significance at $p < 0.001$. Bar with different superscripts (a, b, c, and d) differ from each other.

- | | | | |
|----|--------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------|
| 16 | Figure 5.3.2 | Resettlement of serum FSH, LH and Estradiol in arsenic induced toxicity with co-administration of Vit -B ₁₂ and/ or folic acid. Each bar represents mean \pm SE, N=6. ANOVA followed by two tailed student's t test were used to find out statistical significance at $p < 0.001$. Bar with different superscripts (a, b, c, and d) differ from each other. | 88 |
| 17 | Figure 5.3.3 | SOD and catalase activity shows differential pattern in uterine tissue of the female albino rats. [A] represents SOD activity measured in cell free extracts. Since the higher dose of vitamin B ₁₂ (0.07 μ g) and folic acid (4.0 μ g) alone or in combination in cell free extract depicted more pronounced protection of SOD and catalase activities. Hence, in case of [B], tissue extracts from rats (co-treated with these above comparatively higher doses of these two B vitamins) containing protein in each lane was electrophoresed on 12% and 8.0 % native gel followed by substrate specific development of SOD and catalase bands respectively. [C] Represents densitometric analysis with respect to control. Each bar represents mean \pm SE, N=6. ANOVA followed by two tailed student t test were used to find out statistical significance at * $p < 0.05$; ** $p < 0.01$. | 89 |
| 18 | Figure 5.3.4 | Total peroxidase activities in cell free extract [A] protected better with comparatively higher dose of Vitamin B ₁₂ (0.07 μ g) and folic acid (4.0 μ g). Electrozymogram showing total peroxidase profile [B] using tissue extracted protein and comparatively these two above higher doses of B vitamins considered in the | 90 |

zymogram. The electrophoresis was carried out using conventional 8% polyacrylamide gel. The gels were stained for peroxidase activity using benzidine, glacial acetic acid and H₂O₂. In panel [C] relative densitometric values of peroxidase band from experimental group, as % of the control lane are shown. Each bar represents mean \pm SE, N=6. ANOVA followed by two tailed students t test were used to find out statistical significance at *p<0.01; **p<0.001.

- 19 Figure 5.3.5 Total serum LDH activity illustrated for the extent of cellular damage. [A] is showing enzymatic activity in cell free assay system following dose dependent co-administration of B vitamins in arsenic intoxicated rats. Vitamin B₁₂ (0.07 μ g) and folic acid (4.0 μ g) had comparatively higher effectiveness in the restoration of arsenic induced tissue damage. Hence, in panel [B] these above two doses of vitamins were considered and serum protein in each lane was electrophoresed on agarose gel followed by substrate specific development of LDH bands. Each bar represents mean \pm SE, N=6. ANOVA followed by two tailed student's t test were used to realize statistical significance at *p<0.05; **p<0.01. 91
- 20 Figure 5.3.6 Protective effect (dose dependent) on malondialdehyde (MDA) and conjugated diene (CD) in uterine tissue by co-administration of Vitamin B₁₂ and folic acid in As³⁺ intoxicated rats. Comparatively higher dose of vitamin B₁₂ (0.07 μ g) and folic acid (4.0 μ g) alone or in combination were more effective on the above protection. Each bar represents mean \pm SE, N=6. ANOVA followed by two-tailed student's t test were used *p<0.05; **p<0.01. 92

- 21 Figure 5.3.7 Figure shown (histology of ovary) dose dependent effect of Vitamin-B₁₂ and folic acid by amelioration to reduce the arsenic mediated follicular atresia in ovary of female rat. **92**
- 22 Figure 5.3.8 Uterine tissue was implanted in paraffin, serially sectioned laterally at 5 μm, stained with eosin and haematoxylin (Harris) and observed under a microscope (magnification 340) to study the uterine histoarchitecture (A-H). Secretary nature of uterine bed was lost following arsenic intoxication (panel- B) and gradual and or significant reversal of secretary tubules was observed following the treatment with these B vitamins (panel E-H). Arrow denotes secretary vesicles. Panel distribution: control rat (panel A) or treated with As³⁺-0.4- ppm (panel B) or As³⁺-0.4-ppm+ vitamin B₁₂-0.04μg (panel C) or As³⁺-0.4-ppm plus Folic acid-2.0 μg (panel D) or As³⁺-0.4 ppm+ vitamin B₁₂-0.04μg + Folic acid-2μg (panel E) or As³⁺-0.4 ppm+ vitamin B₁₂-0.07μg (panel F) or As³⁺-0.4- ppm + Folic acid-4.0 μg (panel G) or As³⁺-0.4-ppm+ vitamin B₁₂-0.07 μg + Folic acid-4.0 μg (panel H).Graphical representation denotes fractal dimension of uterine bed was increased in As³⁺ ingested rats. Subsequent co-administration of vitamin B₁₂ and folic acid gradually and dose dependently decreased the fractal dimension from intoxicated state to convalescent state. NB: There is no unit for Fractal dimension, this is ratio. **93**
- 23 Figure 5.3.9 Figure shown the protective effect in uterine tissue architecture by co-administration of Vit-B₁₂ and /or folic acid in arsenic induced female rat. **94**
- 24 Figure 5.3.10 DNA fragmentation in uterus of female rat treated with arsenic. **95**
- A: Vehicle treated control B: AS (0.4

ppm/100g body wt./day). C: AS (0.4ppm/100g body wt. /day) + Vit-B₁₂ (0.04μg /100g body wt./day). D: AS 0.4ppm/100g body wt. /day) + Folic acid (2μg /100g body wt./day). E: AS (0.4ppm/100g body wt./day) + Vit-B₁₂ (0.04μg /100g body wt./day) + Folic acid (2μg/100g body wt./day). F: AS (0.4ppm/ 100g body wt./day) + Vit-B₁₂ (0.07μg /100g body wt./day). G: AS (0.4 ppm/100g body wt./day) + Folic acid (4μg/100g body wt./day). H: AS (0.4ppm/100g body wt./day) + Vit-B₁₂ (0.07μg/100g body wt./day) + Folic acid (4μg /100g body wt./day).

	25	Figure 5.3.11	Protection of DNA breakage in the comet formation is shown in arsenic induced uterine cell by the co- administration of Vit-B ₁₂ and/ or folic acid in different doses.	96
Experiment-IV	26	Figure 5.4.1	Effect of Vit-B ₁₂ and /or folic acid on Serum Total protein (TP), serum glutamate pyruvate transaminase (SGPT), and serum glutamate oxaloacetate transaminase (SGOT), serum alkaline phosphatase (ALP), serum Urea and Creatinine. Each bar represents mean ± SE, N=6. ANOVA followed by two tailed student's t test were used to find out statistical significance at p<0.001. Bar with same superscripts did not differ from each other.	102
	27	Figure 5.4.2	Effect of Vit-B ₁₂ and /or folic acid on CAT (catalase), SOD (superoxidedismutase), MDA (malondialdehyde) and CD (conjugated diene) in hepatic tissue. Each bar represents here is mean ± SE, N=6, p<0.001. ANOVA followed by multiple student's t test. Bar with same superscripts did not differ from each other.	103
		Figure 5.4.3	Effect of Vit-B ₁₂ and /or folic acid on Catalase (CAT), Superoxidedismutase (SOD) in kidney tissue. Each bar	104

28		represents here is mean \pm SE, N=6, p<0.001 and ANOVA followed by multiple student's t test. Bar with same superscripts did not differ from each other.	
29	Figure 5.4.4	Effect of Vit-B ₁₂ and or folic acid on MDA (malondialdehyde) and CD (conjugated diene) in kidney tissue. Each bar represents here is mean \pm SE, N=6, p<0.001. ANOVA followed by multiple student's t test. Bar with same super scripts did not differ from each other.	104
30	Figure 5.4.5	Figure shows that there is no dissimilarity of kidney histoarchitectural by the administration of Vit-B ₁₂ and/or folic acid in comparison with vehicle treated control of female albino rat.	105
31	Figure 5.4.6	Figure shows that there is no disparity of hepatic histoarchitecture by the administration of Vit-B ₁₂ and/or folic acid in comparison with vehicle treated control of female albino rat.	105
32	Figure 5.4.7	DNA fragmentation of uterus in Vit-B ₁₂ and/or folic acid induced female rat. A:Vehicle treated control B: Vit-B ₁₂ (0.07 μ g/100g body wt. /day) C: Folic acid (4 μ g/100g body wt./day) D: Vit-B ₁₂ (0.07 μ g/100g body wt./day + Folic acid (4 μ g /100g body wt./day)	106
33	Figure 5.4.8	There is no Comet formation by DNA fragmentation in uterus by administration of Vit-B ₁₂ and/or folic acid induced female rat.	106
34	Figure 5.4.9	Restoration of Peroxidase activity in Vit-B ₁₂ and/or folic acid in compare with vehicle treated control. But peroxidase activity is more pronounced in Vit-B ₁₂ and folic acid combination group.	107

Experiment-V	35	Figure 5.5.1	Effect of vitamin B ₁₂ and folic acid on arsenic- induced hepatic function markers (mean ± SE; N=6). As compared with control, <i>P</i> <0.001 (ANOVA followed by multiple comparison two-tailed <i>t</i> test). Same superscript did not differ from each other significantly.	111
	36	Figure 5.5.2	Effect of vitamin B ₁₂ and folic acid on lipid profile of arsenic-treated rats (mean ± SE; N = 6). As compared with control, <i>P</i> <0.001 (ANOVA followed by multiple comparison two-tailed <i>t</i> test). Same superscript did not differ from each other significantly.	111
	37	Figure 5.5.3	Effect of vitamin B ₁₂ and folic acid on arsenic-induced oxidative stress in liver of rats (mean ± SE; N = 6). As compared with control, <i>P</i> < 0.001 (ANOVA followed by multiple comparison two-tailed <i>t</i> test). Same superscript did not differ from each other significantly.	112
	38	Figure 5.5.4	Hepatic histoarchitecture (magnification 400×) of female rat treated with arsenic. Control rat (plate A) or treated with sodium arsenite (plate B) or sodium arsenite + vitamin B ₁₂ + folic acid (plate C).	112
	39	Figure 5.5.5	DNA fragmentation in liver of female rat treated with arsenic. Control rat (lanes 1, 2, and 3) or treated with sodium arsenite (lanes 4, 5, and 6) or sodium arsenite + vitamin B ₁₂ + folic acid (lanes 7 and 8).	113
Experiment-VI	40	Figure 5.6.1	Withdrawal effect of selective formulation of co- administration of Vit-B ₁₂ and/or folic acid in arsenic induced female rat's protection of DNA breakage in the comet formation are shown in arsenic induced uterine cell by the co-administration of Vit-B ₁₂ and/ or folic acid in different doses.	118